

If You Vaccinate Your Child, Learn How To Recognize the Signs and Symptoms of Vaccine Reactions.

VACCINE REACTIONS

MOTHER'S DESCRIPTIONS

High Fever (over 103 F°)

"His temperature was 105°. I had to put cool towels on him to bring it down."

Skin (hives, rashes, swelling)

"There was a big, hot, red swollen lump at the site of the shot that stayed for weeks."

High Pitched Screaming

"It was a pain cry, a shrill scream and lasted for hours and nothing would help."

Collapse/Shock

"She turned white with a blue tinge around her mouth and went completely limp."

Excessive Sleepiness

"He passed out and we couldn't wake him to feed or do anything for over 12 hours."

Convulsion

"Her eyes twitched, her chin trembled, her body went rigid and then would shake."

Brain Inflammation

"He just laid in his crib with his eyes wide open then would arch his back and scream and go unconscious. Now he has seizures."

Behavior Changes

"She won't sleep or eat. She throws herself down and screams for no reason. She was sweet and happy and is now out of control. She changed into a totally different child."

Mental/Physical Regression

"My 18 month old son stopped talking and walking after those shots. He developed severe allergies, constant diarrhea, ear infections and was sick all the time."


Blood Disorders

"She got big purple bruises on her skin. The doctor told me she was bleeding because her blood couldn't clot."

Other (such as loss of muscle control, arthritis, paralysis, sudden death)

If your child's health deteriorates after vaccination, your child may be eligible for federal compensation. Vaccine reactions should be reported to the federal Vaccine Adverse Events Reporting System (VAERS) by calling 1-800-822-7967 and to NVIC's Vaccine Reaction Registry at www.NVIC.org.

Brought to you as
a public education service by the



National Vaccine Information Center (NVIC)

204 Mill Street, Suite B-1 | Vienna, VA 22180 | 703-938-0342 | www.NVIC.org

Vaccine Safety Bulletin

Before You Take the Risk, Find Out What It Is.

America and America's Children are in the midst of an epidemic of chronic disease and disability. Today, the Centers for Disease Control admits that **1 American child in 6 is developmentally delayed.**

In the past 25 years, the number of American children with learning disabilities, ADHD, asthma and diabetes **has more than tripled.** Autism increases range from 200 percent to over 5,000 percent in some states.

In the past 25 years, the number of doses of vaccines that pediatricians give American babies and children under age 6 **has more than doubled.**

In 1969, 1 in 7,100 children were diabetic. ▶ **Today 1 in 450 children is diabetic.**

In 1970, autism affected 4 in 10,000 children. ▶ **Today 1 in 175 children is autistic.**


In 1976, there were 796,000 learning disabled children. ▶ **Today there are 3 million learning disabled children.**

In 1979, there were 2 million asthmatic children. ▶ **Today there are 9 million asthmatic children.**

In 1997, there were 1.6 million children with ADHD. ▶ **Today there are 4 million children with ADHD.**

In 1980, the Centers for Disease Control (CDC) and American Academy of Pediatrics (AAP) said children should get 23 doses of 7 vaccines by age six years with the first vaccinations given at 2 months old. ▶ **Today the CDC and AAP tell pediatricians to give children 48 doses of 14 vaccines by age 6 with the first dose given at 12 hours old in the newborn nursery. At 6 months old a child can receive 8 vaccines on a single day. At age 15 months, a child can receive as many as 12 vaccines on a single day.**

Outstanding Question: Is the atypical manipulation of the immune system with more and more vaccines in early life setting some children up for chronic disease and disability? **Is less better?**



National Vaccine Information Center (NVIC)

204 Mill Street, Suite B-1 | Vienna, VA 22180 | 703-938-0342 | www.NVIC.org

48 Doses Of 14 Vaccines Before Age 6?

Before Your Child Is Put At Risk—Learn More About Vaccines.

Birth (12 Hours) Hepatitis B

2 Months

Diphtheria
Tetanus
Pertussis
HIB
Polio
Pneumococcal
Rotavirus
Hepatitis B

4 Months

Diphtheria
Tetanus
Pertussis
HIB
Polio
Pneumococcal
Rotavirus

6 Months

Diphtheria
Tetanus
Pertussis
HIB
Pneumococcal
Rotavirus
Hepatitis B
Flu

15 Months

Diphtheria
Tetanus
Pertussis
HIB
Polio
Measles
Mumps
Rubella
Chicken Pox
Pneumococcal
Flu (plus dose at 2, 3,
and 4 years)
Hepatitis A

5 Years

Diphtheria
Tetanus
Pertussis
Polio
Measles
Mumps
Rubella
Flu
Hepatitis A

Vaccine excipients or ingredients in trace or larger amounts depending on specific vaccine (partial list):
lab altered viruses and bacteria; aluminum; mercury; formaldehyde; phenoxyethanol; glutaraldehyde; sodium chloride; monosodium glutamate (MSG); gelatin; lactose; hydrochloric acid; sorbitol; antibiotics; aluminum sulfate; sodium borate; sodium acetate; hydrogen peroxide; yeast protein; egg albumin; bovine and human serum albumin.

National Vaccine Information Center (NVIC)



204 Mill Street, Suite B-1 | Vienna, VA 22180
703-938-0342 | www.NVIC.org